



Geodesic Mappings of Manifolds with Affine Connection onto the Ricci Symmetric Manifolds

Volodymyr Berezovskii^a, Irena Hinterleitner^b, Josef Mikeš^c

^aUman National University of Horticulture, Dept. of Mathematics and Physics

^bBrno University of Technology, Inst. of Mathematics and Descriptive Geometry

^cPalacky University Olomouc, Dept. of Algebra and Geometry

Abstract. In the present paper we investigate geodesic mappings of manifolds with affine connection onto Ricci symmetric manifolds which are characterized by the covariantly constant Ricci tensor. We obtained a fundamental system for this problem in a form of a system of Cauchy type equations in covariant derivatives depending on no more than $n(n+1)$ real parameters. Analogous results are obtained for geodesic mappings of manifolds with affine connection onto symmetric manifolds.

1. Introduction

The paper is devoted to a further development of Theory of geodesic mappings of manifolds with affine connection. This theory is inspired by T. Levi-Civita [13] where he set and solved in a special system of coordinates the problem of finding Riemannian metrics with common geodesics. It should be noted that it was linked with the study of equations of dynamical mechanical systems.

Later the theory of geodesic mappings was developed in the works of Thomas, Weyl, Cartan, Shirokov, Solodovnikov, Petrov, Sinyukov, Prvanović, Mikeš, etc. [4, 5, 20, 25, 30, 33, 35, 36, 40].

Geodesic mappings play a fundamental role in theoretical mechanics and physics, especially in the general theory of relativity [12, 13, 32, 33], and also in fundamental geometry of special lines structures [29].

Geodesic mappings of more general classes of manifolds (Finsler and generally Einstein) have been studied in the works [22, 31, 43–45].

Geodesic mappings and projective transformations of symmetric, recurrent, Ricci symmetric, Einstein spaces and their generalizations were studied in [7, 14–20, 25–27, 30, 33, 35, 36, 38, 40].

In the fifties of the 20th century Sinyukov [38] studied the geodesic mappings of equiaffine symmetric manifolds with affine connection onto (pseudo-) Riemannian spaces. He proved that these (pseudo-) Riemannian spaces have a constant curvature. This is also valid for recurrent manifolds. These results have been many times repeated, the detailed description see [20, 25, 30, 40].

2010 *Mathematics Subject Classification.* Primary 53B10; Secondary 53B20; 53B30; 53B35; 53B50.

Keywords. manifold with affine connection, symmetric manifold, Ricci symmetric manifold, geodesic mapping, fundamental equations.

Received: 15 February 2017; Revised: 24 February 2017; Accepted: 07 March 2017

Communicated by Mića Stanković

Supported by the project IGA PrF 2017012 Palacky University Olomouc and No. LO1408 "AdMaS UP – Advanced Materials, Structures and Technologies", supported by Ministry of Education, Youth and Sports under the "National Sustainability Programme I" of the Brno University of Technology.

Email addresses: Berez.Volod@rambler.ru (Volodymyr Berezovskii), Hinterleitner.i@fce.vutbr.cz (Irena Hinterleitner), Josef.Mikes@upol.cz (Josef Mikeš)

Mikeš [14, 17–20] has followed up with the study based on the results obtained by Sinyukov. The above mentioned results are also valid for general recurrent spaces. In the paper [7] Hinterleitner and Mikeš proved that the targeted spaces may be the Weyl manifolds.

Similar results are also true for geodesic mappings of non-Einsteinian Ricci symmetric spaces onto (pseudo-) Riemannian spaces [15, 18]. Let us note that geodesic mappings of Einstein spaces have been studied by Mikes [16] and others, see [10, 25, 30].

Sinyukov [39, 40], while studying geodesic mappings between (pseudo-) Riemannian spaces, found out that general solution depends on a finite number of real parameters. This result holds also for geodesic mappings of manifolds with affine connection onto (pseudo-) Riemannian spaces [21], and for the case of geodesic mappings of generalized Finsler spaces onto (pseudo-) Riemannian spaces [22].

In the given study the main equations of geodesic mappings of manifolds A_n with affine connections onto Ricci symmetric manifolds \bar{A}_n were obtained in the form of a system of partial differential equations of Cauchy type. The conditions of their integrability have been found. The numbers of independent real parameters that influence the general solutions of this system have been obtained.

Let us assume that the studied manifolds are simply connected and the dimension of the manifolds is $n \geq 2$. We supposed that geometrical objects are continuous and sufficient differentiable.

2. The Main Concepts of Geodesic Mappings on Manifolds with Affine Connection Theory

Among diffeomorphisms of manifolds geodesic mappings play an important role.

Let us suppose that the manifold A_n with affine connection ∇ admits a diffeomorphism f on the manifold \bar{A}_n with affine connection $\bar{\nabla}$, and these manifolds are related to the common coordinate system (x^1, x^2, \dots, x^n) with respect to f .

We assume that

$$P_{ij}^h(x) = \bar{\Gamma}_{ij}^h(x) - \Gamma_{ij}^h(x) \quad (1)$$

where $\Gamma_{ij}^h(x)$ and $\bar{\Gamma}_{ij}^h(x)$ are components of affine connections ∇ and $\bar{\nabla}$, $P_{ij}^h(x)$ is called a *deformation tensor* of connections with respect to diffeomorphism f .

A curve defined in space A_n is called *geodesic* if its tangent vector is parallel along it. A diffeomorphism $f: A_n \rightarrow \bar{A}_n$ is called a *geodesic mapping* if f maps any geodesic in A_n onto a geodesic in \bar{A}_n .

It is well known [5, 25, 30, 40] that a mapping A_n onto \bar{A}_n is geodesic if and only if in a common coordinate system (x^1, x^2, \dots, x^n) the tensor of deformation has the form

$$P_{ij}^h(x) = \psi_i(x) \delta_j^h + \psi_j(x) \delta_i^h \quad (2)$$

where δ_i^h is the Kronecker symbol and $\psi_i(x)$ are components of a covector.

A geodesic mapping is called *non-trivial* if $\psi_i(x) \neq 0$. It is evident that any arbitrary manifold A_n with affine connection ∇ allows a non-trivial geodesic mapping onto a certain other manifold \bar{A}_n with affine connection $\bar{\nabla}$. Such a suggestion is generally recognized being not true regarding geodesic mapping of (pseudo-) Riemannian spaces onto (pseudo-) Riemannian spaces.

Particularly there were discovered (pseudo-) Riemannian spaces and also spaces with affine connection which do not admit non-trivial geodesic mappings onto (pseudo-) Riemannian spaces [14, 15, 17, 20, 25, 30, 40].

We proved [21] that the main equations of geodesic mappings of A_n onto (pseudo-) Riemannian spaces V_n have a Cauchy type form in covariant derivatives. The main equations of geodesic mappings of equi-affine manifolds on (pseudo-) Riemannian spaces have a linear form. É. Cartan [2] and J.M. Thomas [42] (see [4, p. 105]) demonstrated that any A_n is projectively equivalent with an equi-affine manifold. From this it follows that this system is linear in any case, see [25, p. 282].

3. Geodesic Mappings of Manifolds with Affine Connection onto Ricci Symmetric Manifolds

The manifold A_n with affine connection is called *Ricci symmetric* if the Ricci tensor is covariantly constant. From this it follows that a Ricci symmetric manifold \bar{A}_n is characterized by the condition

$$\bar{R}_{ij|k} \equiv 0 \quad (3)$$

where \bar{R}_{ij} is the Ricci tensor of \bar{A}_n , “|” denotes the covariant derivative with respect to the connection $\bar{\nabla}$ in \bar{A}_n .

Because

$$\bar{R}_{ijk|m}^h = \frac{\partial \bar{R}_{ijk}^h}{\partial x^m} + \Gamma_{\alpha m}^h \bar{R}_{ijk}^\alpha - \Gamma_{im}^\alpha \bar{R}_{\alpha jk}^h - \Gamma_{jm}^\alpha \bar{R}_{i\alpha k}^h - \Gamma_{km}^\alpha \bar{R}_{ij\alpha}^h$$

and considering formula (1) we can write

$$\bar{R}_{ijk|m}^h = \bar{R}_{ijk,m}^h + P_{\alpha m}^h \bar{R}_{ijk}^\alpha - P_{im}^\alpha \bar{R}_{\alpha jk}^h - P_{jm}^\alpha \bar{R}_{i\alpha k}^h - P_{km}^\alpha \bar{R}_{ij\alpha}^h \quad (4)$$

where comma denotes the covariant derivative with respect to the connection ∇ in A_n , \bar{R}_{ijk}^h are components of the Riemann tensor of \bar{A}_n .

After contracting (4) with respect to the indices h and k we get

$$\bar{R}_{ij|m} = \bar{R}_{ij,m} - P_{im}^\alpha \bar{R}_{\alpha j} - P_{jm}^\alpha \bar{R}_{i\alpha}. \quad (5)$$

Further let us suppose that \bar{A}_n is Ricci symmetric. Basing on formula (3) we obtain

$$\bar{R}_{ij,m} = P_{im}^\alpha \bar{R}_{\alpha j} + P_{jm}^\alpha \bar{R}_{i\alpha}. \quad (6)$$

Let us consider geodesic mappings of manifolds A_n with affine connection onto a Ricci symmetric manifold \bar{A}_n . Because the deformation tensor $P_{ij}^h(x)$ of connections has the form (2), than basing on formula (6), we obtain

$$\bar{R}_{ij,m} = 2\psi_m \bar{R}_{ij} + \psi_i \bar{R}_{mj} + \psi_j \bar{R}_{im}. \quad (7)$$

It is known [4, 5, 25, 30, 40] that between the Riemann tensors R_{ijk}^h and \bar{R}_{ijk}^h of the manifolds A_n and \bar{A}_n , respectively, there is a dependence

$$\bar{R}_{ijk}^h = R_{ijk}^h + P_{ik,j}^h - P_{ij,k}^h + P_{ik}^\alpha P_{\alpha j}^h - P_{ij}^\alpha P_{\alpha k}^h. \quad (8)$$

Considering that

$$P_{ij,k}^h(x) = \psi_{i,k}(x) \delta_j^h + \psi_{j,k}(x) \delta_i^h$$

from formula (8) after computation we get

$$\bar{R}_{ijk}^h = R_{ijk}^h - \psi_{i,k} \delta_j^h + \psi_{i,j} \delta_k^h - (\psi_{j,k} - \psi_{k,j}) \delta_i^h + \psi_i \psi_k \delta_j^h - \psi_i \psi_j \delta_k^h. \quad (9)$$

If we contract (9) with respect to the indices h and k we will obtain the following

$$\bar{R}_{ij} = R_{ij} + n \psi_{i,j} - \psi_{j,i} - (n-1) \psi_i \psi_j. \quad (10)$$

After alternation the equation (10) with respect to the indices i and j we can write

$$\bar{R}_{[ij]} = R_{[ij]} + (n+1) (\psi_{i,j} - \psi_{j,i}) \quad (11)$$

where $[ij]$ denotes alternation with respect to the indices i and j .

From equation (11) we find

$$\psi_{i,j} - \psi_{j,i} = \frac{1}{n+1} (\bar{R}_{[ij]} - R_{[ij]}) \quad (12)$$

and from equation (10) considering (12) we have the following formula

$$\psi_{i,j} = \frac{1}{n^2 - 1} [n\bar{R}_{ij} + \bar{R}_{ji} - (nR_{ij} + R_{ji})] + \psi_i \psi_j. \quad (13)$$

It is evident that the equations (7) and (13) in the given manifold A_n represent a closed Cauchy type system with respect to the unknown functions $\bar{R}_{ij}(x)$ and $\psi_i(x)$.

Finally, we obtain the following theorem.

Theorem 3.1. *A manifold A_n with affine connection admits a geodesic mapping onto a Ricci symmetric manifold \bar{A}_n if and only if in A_n exists a solution of a closed system of Cauchy type equations in covariant derivative (7) and (13) with respect to the unknown functions $\bar{R}_{ij}(x)$ and $\psi_i(x)$.*

General solution of a closed system of equations (7) and (13) depends on no more than $n(n+1)$ independent real parameters.

It is evident that the equations (7) and (13) are non linear with respect to the unknown functions $\bar{R}_{ij}(x)$ and $\psi_i(x)$. We obtain the integrability conditions of them.

We differentiate equation (7) covariantly along x^k in the manifold A_n , and then alternate it with respect to the indices m and k . Considering the Ricci identity and formulas (7) and (13) after transformations we can write

$$\begin{aligned} & (n^2 - 1) (\bar{R}_{\alpha j} R_{ikm}^\alpha + \bar{R}_{i\alpha} R_{jkm}^\alpha) - 2(n-1) (\bar{R}_{km} - \bar{R}_{mk}) \bar{R}_{ij} - \\ & n (\bar{R}_{kj} - \bar{R}_{jk}) \bar{R}_{im} - (\bar{R}_{mi} - \bar{R}_{im}) \bar{R}_{kj} - n (\bar{R}_{jm} - \bar{R}_{mj}) \bar{R}_{ki} - \\ & (\bar{R}_{ik} - \bar{R}_{ki}) \bar{R}_{mj} + (nR_{jm} + R_{mj}) \bar{R}_{ik} + (nR_{im} + R_{mi}) \bar{R}_{kj} - \\ & (nR_{ik} + R_{ki}) \bar{R}_{mj} - (nR_{jk} + R_{kj}) \bar{R}_{im} = 0. \end{aligned} \quad (14)$$

Differentiating equation (13) covariantly along x^k in A_n , then alternating with respect to the indices j and k , considering the Ricci identity and formulas (7) and (13), we obtain after transformations

$$\begin{aligned} & (n^2 - 1) \psi_i R_{ijk}^\alpha + n\psi_j R_{ik}^\alpha + \psi_j R_{ki} - n\psi_k R_{ij} - \psi_k R_{ji} + \\ & (n-1) \psi_i (R_{jk} - R_{kj}) = nR_{ik,j} - nR_{ij,k} + R_{ki,j} - R_{ji,k}. \end{aligned} \quad (15)$$

It is evident that equation (15) is linear with respect to the ψ_i (from \bar{R}_{ij} it does not depend).

Equation (14) will be linear with respect to \bar{R}_{ij} (from ψ_i does not depend), if the symmetric manifold A_n ($\bar{R}_{ijk|m}^h = 0$) is also equiaffine.

The manifold A_n is called *equiaffine*, if the Ricci tensor is symmetric, i.e.

$$\bar{R}_{ij} = \bar{R}_{ji}.$$

Thus, the integrability condition of the equations (7) and (13) under geodesic mappings of the manifolds A_n with affine connection on equiaffine symmetric manifolds \bar{A}_n will be linear with respect to the unknown functions $\bar{R}_{ij}(x)$ and $\psi_i(x)$.

4. Geodesic Mappings of Manifolds with Affine Connection onto Symmetric Manifolds

The manifold \bar{A}_n with affine connection $\bar{\nabla}$ is called (locally) *symmetric* if its Riemann tensor is covariantly constant (P.A. Shirokov [37], E. Cartan [2], S. Helgason [6]).

Thus, symmetric manifolds \bar{A}_n are characterized by the condition

$$\bar{R}_{ijk|m}^h = 0. \quad (16)$$

It is evident that symmetric manifolds are also simultaneously Ricci symmetric manifolds. On the other hand, symmetric manifolds are semisymmetric. In 1950 these spaces were introduced by Sinyukov [38], they are characterized by the following condition

$$\bar{R}_{ijk|lm}^h - \bar{R}_{ijk|ml}^h = 0.$$

Geodesic mappings of semisymmetric spaces were studied by Sinyukov [38, 40] and Mikeš [14, 15, 17–20, 23, 25, 26, 28, 30, 40].

Let us consider a geodesic mapping f of the manifold A_n with affine connection ∇ onto a symmetric manifold \bar{A}_n with affine connection $\bar{\nabla}$.

Equations (7) and (13) are valid for the above mentioned geodesic mapping, because, as we have mentioned, the symmetric manifold \bar{A}_n is also Ricci symmetric.

Due to the condition (16) from equation (4) we get

$$\bar{R}_{ijk,m}^h = P_{im}^\alpha \bar{R}_{\alpha jk}^h + P_{jm}^\alpha \bar{R}_{i\alpha k}^h + P_{km}^\alpha \bar{R}_{ij\alpha}^h - P_{\alpha m}^h \bar{R}_{ijk}^\alpha \quad (17)$$

Taking into consideration that the deformation tensor $P_{ij}^h(x)$ has the structure (2), from (17) we obtain the following equation

$$\bar{R}_{ijk,m}^h = 2\psi_m \bar{R}_{ijk}^h + \psi_i \bar{R}_{mjk}^h + \psi_j \bar{R}_{imk}^h + \psi_k \bar{R}_{ijm}^h - \delta_m^h \psi_\alpha \bar{R}_{ijk}^\alpha. \quad (18)$$

Contracting (18) on the indices h and k , we obtain equation (7). This means that equation (7) is a consequence of equation (18).

It is evident that equations (13) and (18) in the given manifold A_n are a Cauchy type closed system of differential equations regarding the unknown functions $\bar{R}_{ijk}^h(x)$ and $\psi_i(x)$ that, naturally, should satisfy moreover the final conditions of algebraic character

$$\bar{R}_{i(jk)}^h(x) = 0, \quad \bar{R}_{(ijk)}^h(x) = 0 \quad (19)$$

where brackets mean the symmetrization of the given indices without division.

From the above mentioned we prove the following theorem.

Theorem 4.1. *The manifold A_n with affine connection admits a geodesic mapping onto a symmetric manifold \bar{A}_n if and only if in A_n exists a solution of a mixed system of Cauchy type equations in covariant derivative (13), (18) and (19) with respect to the unknown functions $\bar{R}_{ijk}^h(x)$ and $\psi_i(x)$.*

It is evident that the general solution of the mixed closed system of Cauchy type equations depends no more that from

$$\frac{1}{2} n^3(n-1) + n$$

independent parameters.

References

- [1] V. Berezovskii, J. Mikeš, P. Peška, Geodesic mappings of manifolds with affine connection onto symmetric manifolds, Proc. 11th Int. Conf. on Geometry, Integrability and Quantization, (Sofia: Avangard Prima, 2017), 99–104.
- [2] É. Cartan, Sur les variétés à connexion projective, S. M. F. Bull. 52 (1924), 205–241.
- [3] H. Chudá, J. Mikeš, Conformally geodesic mappings satisfying a certain initial condition, Arch. Math. (Brno) 47 (2011), 389–394.
- [4] L. P. Eisenhart, Non-Riemannian geometry, AMS, Providence, 1990.
- [5] L. P. Eisenhart, Riemannian geometry, Princeton Univ. Press, 1997.
- [6] S. Helgason, Differential geometry, Lie groups, and symmetric spaces, AMS, 1978.
- [7] I. Hinterleitner, J. Mikeš, Geodesic mappings onto Weyl manifolds, J. Appl. Math. 2 (2009), 125–133.
- [8] I. Hinterleitner, J. Mikeš, Fundamental equations of geodesic mappings and their generalizations, J. Math. Sci. 174 (2011), 537–554.
- [9] I. Hinterleitner, J. Mikeš, Projective equivalence and manifolds with equiaffine connection, J. Math. Sci. 177 (2011), 546–550.
- [10] I. Hinterleitner, J. Mikeš, Geodesic mappings and Einstein spaces, in Geom. meth. phys., Birkhäuser/Springer, Trends Math. 19 (2013), 331–335.
- [11] I. Hinterleitner, J. Mikeš, Geodesic mappings and differentiability of metrics, affine and projective connections, Filomat 29 (2015) 1245–1249.
- [12] S. Ivanov, M. Zlatanović, Connections on a non-symmetric (generalized) Riemannian manifold and gravity, Classical Quantum Gravity 33 (2016), no. 7, 075016, 23 pp.
- [13] T. Levi-Civita, Sulle trasformazioni delle equazioni dinamiche, Ann. Mat. Milano 24 (1896), 255–300.
- [14] J. Mikeš, Geodesic mappings of semisymmetric Riemannian spaces, Arch. VINITI (Moscow) 3924-76 (1976), 1–19.
- [15] J. Mikeš, On geodesic mappings of 2-Ricci symmetric Riemannian spaces. Math. Notes 28, 622–624, 1981. Transl. from Mat. Zametki 28, 313–317, 1980.
- [16] J. Mikeš, On geodesic mappings of Einstein spaces. Math. Notes 28, 922–923, 1981. Transl. from Mat. Zametki 28, 935–938, 1980.
- [17] J. Mikeš, Geodesic mappings of m -symmetric and generalized semisymmetric spaces, Russian Math. (Iz. VUZ) 36 (1992), 38–42. Transl. from Izv. VUZ, Mat. 8 (1992), 42–46.
- [18] Mikeš J. On geodesic and holomorphic-projective mappings of generalized m -recurrent Riemannian spaces. Sib. Mat. Zh. 33:5, 215, 1992.
- [19] J. Mikeš, Geodesic mappings onto semisymmetric spaces. Russ. Math. 38:2, 35–41, 1994. Transl. from Izv. VUZ, Mat. 381:2, 37–43, 1994.
- [20] J. Mikeš, Geodesic mappings of affine-connected and Riemannian spaces. J. Math. Sci. (New York) 78:3, 311–333, 1996.
- [21] J. Mikeš, V. Berezovski, Geodesic mappings of affine-connected spaces onto Riemannian spaces, Amsterdam: North-Holland. Colloq. Math. Soc. J. Bolyai, 56. Diff. Geom. Eger Hungary, 491–494, 1989.
- [22] J. Mikeš, S. Bácsó, V. Berezovskii, Geodesic mappings of weakly Berwald spaces and Berwald spaces onto Riemannian spaces, Int. J. Pure Appl. Math. 45 (2008), no. 3, 413–418.
- [23] J. Mikeš, V.E. Berezovski, E.S. Stepanova, H. Chudá, Geodesic mappings and their generalizations, J. Math. Sci. 217 (2016), 607–623.
- [24] J. Mikeš, H. Chudá, I. Hinterleitner, Conformal holomorphically projective mappings of almost Hermitian manifolds with a certain initial condition, Int. J. Geom. Methods Mod. Phys. 11 (2014), 1450 044, 8.
- [25] J. Mikeš et al., Differential geometry of special mappings, Palacky Univ. Press, Olomouc, 2015.
- [26] J. Mikeš, Ž. Radulović, On geodesic and holomorphically projective mappings of generalized recurrent spaces. Publ. Inst. Math. 59(73), 153–160, 1996.
- [27] J. Mikeš, Ž. Radulović, On projective transformations of Riemannian spaces with harmonic curvature. New developments in diff. geom., Budapest 1996. Kluwer Acad. Publ. 279–283, 1999.
- [28] J. Mikeš, Ž. Radulović, M. Haddad, Geodesic and holomorphically projective mappings of m -pseudo- and m -quasisymmetric Riemannian spaces. Russ. Math. 40:10, 28–32, 1996.
- [29] J. Mikeš, K. Strambach, Differentiable structures on elementary geometries, Results Math. 53 (2009), no. 1–2, 153–172.
- [30] J. Mikeš, A. Vanžurová, I. Hinterleitner, Geodesic mappings and some generalizations, Palacky Univ. Press, Olomouc, 2009.
- [31] M.S. Najdanović, M.Lj. Zlatanović, I. Hinterleitner, Conformal and geodesic mappings of generalized equidistant spaces, Publ. Inst. Math. (Beograd) (N.S.) 98(112) (2015), 71–84.
- [32] A.Z. Petrov, On geodesic mappings of Einstein spaces, Izv. VUZ, Mat. 21, 130–136, 1961.
- [33] A.Z. Petrov, New methods in the general theory of relativity, Nauka, Moscow, 1966.
- [34] M. Prvanović, Projective and conformal transformations in recurrent and Ricci recurrent Riemannian spaces. Tensor 12:13, 219–226, 1962.
- [35] M. Prvanović, Projektive Geometrie, (Serbo-Croat) Nauchna knjiga, Beograd, 1968.
- [36] Zh. Radulovich, J. Mikeš, M.L. Gavril'chenko, Geodesic mappings and deformations of Riemannian spaces. CID, Podgorica; OGU, Odessa, 1997.
- [37] P.A. Shirokov, Selected investigations on geometry, Kazan Univ. press, 1966.
- [38] N.S. Sinyukov, On geodesic mappings of Riemannian manifolds onto symmetric spaces, Dokl. Akad. Nauk SSSR 98, 21–23, 1954.
- [39] N.S. Sinyukov, A contribution to the theory of geodesic mapping of Riemannian spaces. Sov. Math., Dokl. 7, 1004–1006, 1966. Transl. from Dokl. Akad. Nauk SSSR 169, 770–772, 1966. Corr. 7:6, v–vi, 1966.
- [40] N.S. Sinyukov, Geodesic mappings of Riemannian spaces, Nauka, Moscow, 1979.
- [41] M.S. Stanković, S.M. Minčić, New special geodesic mappings of generalized Riemannian spaces, Publ. Inst. Math., 67 (2000), 92–102.
- [42] J.M. Thomas, Asymmetric displacement of a vector, Trans. AMS 28 (1926), 658–670.

- [43] M.Lj. Zlatanović, I. Hinterleitner, M. Najdanović, Geodesic mapping onto Kählerian spaces of the first kind, Czechosl. Math. J. 64(139) (2014), no. 4, 1113–1122.
- [44] M.Lj. Zlatanović, Lj.S. Velimirović, M.S. Stanković, Necessary and sufficient conditions for equitorsion geodesic mapping, J. Math. Anal. Appl. 435 (2016), 578–592.
- [45] M.Lj. Zlatanović, V. Stanković, Geodesic mapping onto Kählerian space of the third kind, J. Math. Anal. Appl. 450 (2017), no. 1, 480–489.