

Almost geodesic mappings of affinely connected spaces that preserve the Riemannian curvature*

Vladimir Berezovski^a, Sándor Bácsó^b, Josef Mikeš^c

^aUman National University of Horticulture, Ukraine
berez.volod@rambler.ru

^bDebrecen University, Hungary
bacsos@unideb.hu

^cPalacky University Olomouc, Czech Republic
josef.mikes@upol.cz

Submitted July 22, 2015 — Accepted September 29, 2015

Abstract

In the present paper the authors give some conditions preserved Riemannian curvature tensor with respect to almost geodesic mappings of affinely connected spaces. It is noteworthy that these conditions are valid for other types of mappings. For the almost geodesic mappings of first type, when the Riemannian curvature tensor is invariant, the authors deduce a differential equations system of Cauchy type.

In addition the authors investigate almost geodesic mappings of first type, where the Weyl tensor of projective curvature is invariant and Riemannian tensor is not invariant.

Keywords: almost geodesic mapping, Riemannian curvature tensor

MSC: 53B05

*The paper was supported by the grant IGA Faculty of Science 2014016 Mathematical Structures of the Palacky University

1. Introduction

Several works [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24] have been devoted to study almost geodesic mappings. These mappings are generalization of geodesic and quasigeodesic mappings, see [11, 12, 13, 17].

The basic concepts of almost geodesic curve and almost geodesic mapping of affinely connected spaces are introduced in paper [15] and included in the monographs [17, p. 156], [12, p. 457] and surveys [18, 4, 8, 10].

Definition 1.1. A curve $x(t)$ in an affinely connected space A_n is called an *almost geodesic curve* if there exists a plane $\tau(t)$ in every tangent space of the curve $x(t)$ such that:

- (1) $\tau(t)$ are parallel translated along $x(t)$, and
- (2) the tangent vector $\dot{x}(t)$ of the curve lies in $\tau(t)$.

Definition 1.2. A diffeomorphism $f: A_n \rightarrow \bar{A}_n$ is called *almost geodesic mapping*, if under f any geodesic curve of A_n coincides with an almost geodesic curve of $\bar{A}_n$.

Theorem 1.3. *Diffeomorphism $f: A_n \rightarrow \bar{A}_n$ is almost geodesic mapping if and only if the deformation tensor of the affine connections $P_{ij}^h(x) \equiv \bar{\Gamma}_{ij}^h(x) - \Gamma_{ij}^h(x)$ satisfies for any vector λ^h the following conditions:*

$$A_{\alpha\beta\gamma}^h \lambda^\alpha \lambda^\beta \lambda^\gamma = a P_{\alpha\beta}^h \lambda^\alpha \lambda^\beta + b \lambda^h$$

where

$$A_{ijk}^h = P_{ij,k}^h + P_{ij}^\alpha P_{k\alpha}^h, \quad (1.1)$$

Γ_{ij}^h ($\bar{\Gamma}_{ij}^h$) are objects of affine connections of spaces A_n ($\bar{A}_n$) respectively, a and b are some functions depend on x^h and λ^h and $x = (x^1, x^2, \dots, x^n)$ is a common system of coordinates. The symbol “,” means covariant derivation with respect to A_n .

Three types of almost geodesic mapping was discovered by Sinyukov [15, 16, 17, 18], he called them π_1 , π_2 and π_3 . In [2] it was proved that another almost geodesic mapping, if $n > 5$ does not exist.

Almost geodesic mapping π_1 is characterized by the following conditions:

$$A_{(ijk)}^h = \delta_{(i}^h a_{jk)} + b_{(i} P_{jk)}^h,$$

where a_{ij} is a symmetric tensor, b_i is a vector, and the symbol (ijk) means symmetrization without division for the indices i, j, k .

2. Mappings of affinely connected spaces that preserve the Riemann curvature tensor

If we give a diffeomorphism $f: A_n \rightarrow \bar{A}_n$, then the relation between Riemann curvature tensors R_{ijk}^h and $\bar{R}_{ijk}^h$ of A_n and $\bar{A}_n$ is the following [18, p. 78], [11, p. 86], [12, p. 184]:

$$\bar{R}_{ijk}^h = R_{ijk}^h + P_{i[k,j]}^h + P_{i[k}^\alpha P_{j]\alpha}^h, \quad (2.1)$$

where the symbol $[kj]$ denotes the alternation for the indices k and j .

Using of (1.1) and (2.1) we have

Theorem 2.1. *A mapping preserves the Riemann curvature tensor if and only if it satisfies the condition*

$$A_{ijk}^h = A_{ikj}^h, \quad (2.2)$$

that is, the tensor A_{ijk}^h is to be symmetric in the indices j and k .

If the Riemann curvature tensor is preserved by the mapping, then, of course, Ricci tensor $R_{ij} = R_{i\alpha j}^\alpha$ and Weyl tensor of projective curvature

$$W_{ijk}^h = R_{ikj}^h - \frac{1}{n+1} \delta_i^h R_{[jk]} + \frac{1}{n^2-1} [(nR_{ij} + R_{ji})\delta_k^h - (nR_{ik} + R_{ki})\delta_j^h] \quad (2.3)$$

also are invariant under this mapping.

The condition of Theorem 1.3 is sufficient condition for preserving the Ricci tensor and Weyl tensor of projective curvature, but it is not necessary. Further on we give an example.

3. Special almost geodesic mappings of first type which preserve Riemannian tensor

Let be a mapping given between affinely connected spaces A_n and $\bar{A}_n$, which satisfies the condition:

$$P_{ij,k}^h + P_{ik,j}^h = -P_{ij}^\alpha P_{\alpha k}^h - P_{ik}^\alpha P_{\alpha j}^h + \delta_{(i}^h a_{jk)} \quad (3.1)$$

This mapping is a special case of almost geodesic mapping of first type.

Alternating equation (3.1) in i and j , we get

$$P_{ik,j}^h - P_{jk,i}^h = -P_{ik}^\alpha P_{\alpha j}^h + P_{jk}^\alpha P_{\alpha i}^h. \quad (3.2)$$

At now in (3.2), we exchange the indices i and k , we obtain

$$P_{ik,j}^h - P_{ji,k}^h = -P_{ik}^\alpha P_{\alpha j}^h + P_{ji}^\alpha P_{\alpha k}^h. \quad (3.3)$$

If we subtract equation (3.3) from equation (3.1), we have

$$2P_{ij,k}^h = -2P_{ij}^\alpha P_{\alpha k}^h + \delta_{(i}^h a_{jk)},$$

that is,

$$P_{ij,k}^h + P_{ij}^\alpha P_{\alpha k}^h = \delta_{(i}^h \tilde{a}_{jk)}, \quad (3.4)$$

where $\tilde{a}_{ij} = \frac{1}{2}a_{ij}$.

In this case the tensor $A_{ijk}^h = \delta_{(i}^h \tilde{a}_{jk)}$ is symmetric in indices j and k .

Using of Theorem 1.3 we have

Theorem 3.1. *The almost geodesic mapping (determined by (3.1)) preserves the Riemann curvature tensor R_{ijk}^h .*

If Riemann curvature tensor vanishes in an affine space, then we have the following

Theorem 3.2. *If an affine space A_n admits an almost geodesic mapping (determined by (3.1)) into $\bar{A}_n$, then $\bar{A}_n$ is also an affine space.*

So affinely spaces are closed under almost geodesic mapping (determined by (3.1)).

From equation (3.1) we obtained the equation (3.4). Equation (3.4) is a system of Cauchy type for deformation tensor. We can find it's integrability conditions.

We differentiate covariantly equation (3.4) by x^m , further on, we change the indices k and m , using of Ricci identities we have

$$\delta_i^h \tilde{a}_{j[k,m]} + \delta_j^h \tilde{a}_{i[k,m]} + \delta_{[k}^h \tilde{a}_{ij,lm]} = P_{ij}^\alpha R_{\alpha km}^h + P_{\alpha(j}^h R_{i)km}^\alpha + a_{j[m} \tilde{a}_{k]i}^h + \tilde{a}_{i[m} \tilde{a}_{k]j}^h. \quad (3.5)$$

After transvecting of integrability conditions (3.5) by indices h and m we obtain

$$\begin{aligned} \tilde{a}_{jk,i} + \tilde{a}_{ik,j} - (n+1)\tilde{a}_{ij,k} &= -P_{ij}^\alpha R_{\alpha k} + P_{\alpha j}^\beta R_{ik\beta}^\alpha + P_{\alpha i}^\beta R_{jk\beta}^\alpha \\ &+ \tilde{a}_{j\alpha} P_{ki}^\alpha - \tilde{a}_{jk} P_{\alpha i}^\alpha + \tilde{a}_{i\alpha} P_{jk}^\alpha - \tilde{a}_{ik} P_{j\alpha}^\alpha. \end{aligned} \quad (3.6)$$

Alternating equation (3.6) in k and j , we obtain

$$\begin{aligned} \tilde{a}_{ij,k} &= \tilde{a}_{ik,j} + \frac{1}{n+2}(-P_{ij}^\alpha R_{\alpha k} + P_{ik}^\alpha R_{\alpha j} - P_{\alpha j}^\beta R_{ik\beta}^\alpha + P_{\alpha k}^\beta R_{ij\beta}^\alpha - \\ &P_{\alpha i}^\beta R_{jk\beta}^\alpha + P_{\alpha k}^\beta R_{ij\beta}^\alpha - \tilde{a}_{j\alpha} P_{ki}^\alpha + \tilde{a}_{k\alpha} P_{ij}^\alpha + \tilde{a}_{ik} P_{j\alpha}^\alpha - \tilde{a}_{ij} P_{k\alpha}^\alpha). \end{aligned} \quad (3.7)$$

In equation we exchange the indices k and i , we get

$$\begin{aligned} \tilde{a}_{kj,i} &= \tilde{a}_{ik,j} + \frac{1}{n+2}(-P_{kj}^\alpha R_{\alpha i} + P_{ki}^\alpha R_{\alpha j} - P_{\alpha j}^\beta R_{ki\beta}^\alpha + P_{\alpha i}^\beta R_{kj\beta}^\alpha - \\ &P_{\alpha k}^\beta R_{ji\beta}^\alpha + P_{\alpha k}^\beta R_{ij\beta}^\alpha - \tilde{a}_{j\alpha} P_{ik}^\alpha + \tilde{a}_{i\alpha} P_{kj}^\alpha + \tilde{a}_{ki} P_{j\alpha}^\alpha - \tilde{a}_{kj} P_{i\alpha}^\alpha). \end{aligned} \quad (3.8)$$

Substituting equation (3.7) and (3.8) into (3.6), we have

$$\begin{aligned} \tilde{a}_{ik,j} &= \frac{1}{(n-1)(n+2)}(-n(P_{ik}^\alpha R_{\alpha j} + P_{\alpha(k}^\beta R_{i)j\beta}^\alpha)) - R_{\alpha(k} P_{i)j}^\alpha - P_{\alpha j}^\beta R_{(ik)\beta}^\alpha - \\ &P_{\alpha(i}^\beta R_{j)k\beta}^\alpha + (n+1)(\tilde{a}_{j(i} P_{k)\alpha}^\alpha - \tilde{a}_{\alpha(i} P_{k)j}^\alpha) + 2(\tilde{a}_{ik} P_{j\alpha}^\alpha - \tilde{a}_{j\alpha} P_{ik}^\alpha). \end{aligned} \quad (3.9)$$

Equation (3.4) and (3.9) are a system of Cauchy type for the function $P_{ij}^h(x)$ and $\tilde{a}_{ij}(x)$, which satisfy the following

$$P_{ij}^h(x) = P_{ji}^h(x), \quad \tilde{a}_{ij}(x) = \tilde{a}_{ji}(x). \quad (3.10)$$

So is proved the

Theorem 3.3. *An affinely connected space A_n admits an almost geodesic mapping (determined by (3.1)) into an affinely connected space $\bar{A}_n$ if and only if in A_n exist solution functions $P_{ij}^h(x)$ and $\bar{a}_{ij}(x)$ for equation system of Cauchy type (3.4), (3.9) and (3.10).*

4. Special almost geodesic mappings of first type which preserve Weyl tensor of projective curvature but does not preserve Riemann curvature tensor

Let be the $f: \bar{A}_n \rightarrow \bar{A}_n$ mapping given, which satisfies the following condition

$$P_{ij,k}^h + P_{ij}^\alpha P_{\alpha k}^h = \delta_k^h a_{ij}, \quad (4.1)$$

where a_{ij} is a symmetric tensor.

It is well known, that this above mentioned mapping is an almost geodesic mapping of first type.

The tensor A_{ijk}^h on the basis of (4.1) is equal to $\delta_k^h a_{ij}$. If the tensor $a_{ij} \neq 0$, then the tensor A_{ijk}^h is not symmetric in indices j and k . So, in general the mapping (determined by (4.1)) does not preserve the Riemannian curvature tensor.

Using of (2.1) and (4.1) we get

$$\bar{R}_{ijk}^h = R_{ijk}^h - a_{i[j} \delta_{k]}^h. \quad (4.2)$$

It is easy to see, that after transvecting (4.2) in indices h and k , we have

$$a_{ij} = \frac{1}{n-1} (\bar{R}_{ij} - R_{ij}).$$

From the last formulae, symmetry a_{ij} , (2.3) and (4.2) we get

$$\bar{W}_{ij} = W_{ij}, \quad \bar{\tilde{W}}_{ijk}^h = \tilde{W}_{ijk}^h, \quad \text{and} \quad \bar{W}_{ijk}^h = W_{ijk}^h$$

where

$$W_{ij} = R_{ij} - R_{ji} \quad \text{and} \quad \tilde{W}_{ijk}^h = R_{ijk}^h + \frac{1}{n-1} R_{i[j} \delta_{k]}^h.$$

The W_{ij} and $\bar{W}_{ij}$ are tensors of type $\binom{0}{2}$ in the space A_n and $\bar{A}_n$ respectively. The $\tilde{W}_{ijk}^h$ and $\bar{\tilde{W}}_{ijk}^h$ are tensors of type $\binom{1}{3}$ in the space A_n and $\bar{A}_n$ respectively. The W_{ijk}^h and $\bar{W}_{ijk}^h$ are Weyl tensors of projective curvature of A_n and $\bar{A}_n$ respectively.

Finally we obtain

Theorem 4.1. *Tensors W_{ij} , $\tilde{W}_{ijk}^h$ and W_{ijk}^h are invariants under almost geodesic mapping (determined by (4.1)).*

From Theorem 3.3 follows

Theorem 4.2. *If a projective-euclidean or equiaffinely space A_n admits almost geodesic mapping (determined by (4.1)) into an affinely connected space $\bar{A}_n$, then $\bar{A}_n$ is a projective-euclidean or equiaffinely space respectively.*

The proof of Theorem 4.1 follows from facts, that the Weyl tensor vanishes in projective-euclidean space, and the tensor is equal to zero in equiaffinely space.

So, using of Theorem 4.1, we obtain, that the projective-euclidean and equiaffinely spaces are closed sets under almost geodesic mapping (determined by (4.1)).

For almost geodesic mapping of first type, which determined by (4.1), the tensor A_{ijk}^h is equal to $\delta_i^h a_{ij}$. If $a_{ij} \neq 0$, then the A_{ijk}^h tensor is not symmetric in indices j and k .

So, the mapping (determined by (4.1)) does not preserve the Riemann curvature tensor, but the Weyl tensor is invariant object.

Consider the equations (4.1) as a system of Cauchy-type for unknown P_{ij}^h , find it's integrability condition. At first, we differentiate covariantly equation (4.1) in x^m , further on we alternate it in indices k and m . After transvecting integrability condition of equation (4.1) in indices h and m , we obtain

$$(n-1)a_{ij,k} = P_{ij}^\alpha R_{\alpha k} - P_{\alpha(i} R_{j)\beta k}^\alpha - (n-1)P_{ij}^\alpha a_{\alpha k} \quad (4.3)$$

So equation (4.1) and (4.3) in a space A_n give a system of Cauchy type for unknown functions $P_{ij}^h(x)$ and $a_{ij}(x)$, which satisfies algebraic conditions

$$P_{ij}^h(x) = P_{ji}^h(x), \quad a_{ij}(x) = a_{ji}(x) \quad (4.4)$$

Therefore

Theorem 4.3. *An affinely connected space admits almost geodesic mapping of first type (determined by (4.1)) into an affinely connected space $\bar{A}_n$ if and only if in A_n there exist solution $P_{ij}^h(x)$ and $a_{ij}(x)$ for system of Cauchy type equations (4.1), (4.3) and (4.4).*

References

- [1] AMINOVA, A. V., Group of almost projective motions of spaces with affine connection, *Sov. Math.*, 23:4 (1979) 71–74. Transl. from *Izv. vuzov, Mat.*, 4 (1979) 71–75.
- [2] BEREZOVSKI, V. E., MIKEŠ, J., On a classification of almost geodesic mappings of affine connection spaces, *Acta Univ. Palacki. Olomuc., Math.*, 35 (1996) 21–24.
- [3] BEREZOVSKI, V. E., MIKEŠ, J., Almost geodesic mappings of type π_1 onto generalized Ricci-symmetric manifolds, *Uch. zap. Kazan. Univ. Ser. Fiz.-Math.*, 151:4 (2009) 9–14.
- [4] BEREZOVSKI, V. E., MIKEŠ, J., Almost geodesic mappings of spaces with affine connection. *J. Math. Sci. New York*, 2015. Transl. from *Itogi nauk, RAN VINITI*, 126 (2013) 62–95.

- [5] BEREZOVSKI, V. E., MIKEŠ, J., On canonical almost geodesic mappings of the first type of affinely connected spaces, *Russ. Math.*, 58:2 (2014) 1–5. Transl. from *Izv. vuzov, Mat.*, 2 (2014) 3–8.
- [6] BEREZOVSKI, V. E., MIKEŠ, J., VANŽUROVÁ, A., Canonical almost geodesic mappings of type π_1 onto pseudo-Riemannian manifolds, *DGA Proc. Conf., World Sci.*, (2008) 65–76.
- [7] BEREZOVSKI, V. E., MIKEŠ, J., VANŽUROVÁ, A., Almost geodesic mappings onto generalized Ricci-symmetric manifolds, *Acta Math. Acad. Paed. Nyházi. (N.S.)* 26:2 (2010) 221–230.
- [8] BEREZOVSKI, V. E., MIKEŠ, J., VANŽUROVÁ, A., Fundamental PDE's of the canonical almost geodesic mappings of type $\tilde{\pi}_1$, *Bull. Malays. Math. Sci. Soc.*, 37:3 (2014) 647–659.
- [9] FORMELLA, S., A note on geodesic and almost geodesic mappings of homogeneous Riemannian manifolds, *Opusc. Math.*, 25:2 (2005) 181–187.
- [10] MIKEŠ, J., Holomorphically projective mappings and their generalizations, *J. Math. Sci. (New York)*, 89:3 (1998) 1334–1353. Transl. from *Itogi Nauki Tekh., Ser. Sovrem. Mat. Prilozh., Temat. Obz.*, 30 (2002) 258–289.
- [11] MIKEŠ, J., HINTERLEITNER, I., VANŽUROVÁ, A., Geodesic mappings and some generalizations, *Palacky Univ. Press, Olomouc*, (2009).
- [12] MIKEŠ, J., AT AL., Differential geometry of special mappings, *Palacky Univ. Press, Olomouc*, (2015).
- [13] PETROV, A. Z., Modeling of the paths of test particles in gravitation theory, *Gravit. and the Theory of Relativity*, 4-5 (1968) 7–21.
- [14] PETROVIĆ, M. Z., STANKOVIĆ, M. S., Special almost geodesic mappings of the first type of non-symmetric affine connection spaces, *Bull. Malays. Math. Sci. Soc.*, DOI 10.1007/s40840-015-0118-0 (2015).
- [15] SINYUKOV, N. S., Almost geodesic mappings of affinely connected and Riemannian spaces, *Sov. Math., Dokl.*, 4 (1963) 1086–1088. Transl. from *Dokl. Akad. Nauk SSSR*, 151 (1963) 781–782.
- [16] SINYUKOV, N. S., Almost-geodesic mappings of affinely-connected spaces and e-structures, *Math. Notes*, 7 (1970) 272–278.
- [17] SINYUKOV, N. S., Geodesic mappings of Riemannian spaces, *Nauka, Moscow*, (1979).
- [18] SINYUKOV, N. S., Almost-geodesic mappings of affinely connected and Riemann spaces, *J. Sov. Math.*, 25 (1984) 1235–1249. Transl. from *Itogi Nauki Tekh., Ser. Probl. Geom.*, 13 (1982) 3–26.
- [19] SOBCHUK, V. S., Almost geodesic mappings of Riemann spaces onto symmetric Riemann spaces, *Math. Notes*, 17 (1975) 450–454.
- [20] SOBCHUK, V. S., Interior almost geodesic mappings, *Sov. Math.*, 33:5 (1989) 97–101. Transl. from *Izv. vuzov, Mat.*, 324:5 (1989) 62–64.
- [21] SOBCHUK, V. S., MIKEŠ, J., POKORNÁ, O., On almost geodesic mappings π_2 between semisymmetric Riemannian spaces, *Novi Sad J. Math.*, 29:3 (1999) 309–312.
- [22] STANKOVIĆ, M. S., On canonic almost geodesic mappings of the second type of affine spaces, *Filomat*, **13** (1999) 105–144.

- [23] VAVŘÍKOVÁ, H., MIKEŠ, J., POKORNÁ, O., STARKO, G., On fundamental equations of almost geodesic mappings of type $\pi_2(e)$, *Russ. Math.*, 51:1 (2007) 8–12. Transl. from *Izv. vuzov, Mat.*, 1 (2007) 10–15.
- [24] YABLONSKAYA, N. V., Special groups of almost geodesic transformations of spaces with affine connection, *Sov. Math.*, 30:1 (1986) 105–108. Transl. from *Izv. vuzov, Mat.*, 284:1 (1986) 78–80.